

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Postnatal Vitamin A Supplementation in Developing Countries: An Intervention Whose Time Has Come?

Steven A. Abrams and David C. Hilmers

Pediatrics 2008;122:180-181

DOI: 10.1542/peds.2008-0455

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://www.pediatrics.org/cgi/content/full/122/1/180>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2008 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Postnatal Vitamin A Supplementation in Developing Countries: An Intervention Whose Time Has Come?

Steven A. Abrams, MD, David C. Hilmers, MD

Department of Pediatrics, Baylor College of Medicine, Houston, Texas

The authors have indicated they have no financial relationships relevant to this article to disclose.

IN THIS MONTH'S *Pediatrics* Electronic Pages, Klemm et al¹ have provided new information that demonstrates a mortality benefit to neonatal vitamin A supplementation (VAS) in a developing country. Although nearly 100 years have passed since vitamin A was discovered, and more than 60 years since it was first synthesized, the precise mechanisms of its beneficial effects as a supplement remain to be elucidated. Beginning with trials reported in the 1980s,² VAS in children ≥ 6 months of age has been accepted as a cost-effective intervention that is associated with a significant decrease in all-cause mortality in children aged 6 to 59 months in areas of highly prevalent vitamin A deficiency.³ It is estimated that in 1999 VAS prevented 242 000 childhood deaths worldwide at an incremental cost of \$64 per death averted and \$0.02 per dose.⁴ Few significant adverse effects have been reported. VAS has been shown to be effective in reducing chronic lung disease in very low birth weight infants.⁵ However, not all studies have been positive; trials have shown no benefit of VAS in infants aged 1 to 5 months.⁶

The evidence for postnatal VAS in neonates born near term has been mixed. Klemm et al administered 50 000 IU of vitamin A to infants within the first month of life in Bangladesh, a large majority of whom were born at home.¹ The relative risk of all-cause mortality (0.70–1.00) was reduced by 15% ($P = .045$) for those who received VAS versus placebo, and the trial was stopped early after the survival benefit became apparent. This positive outcome is consistent with 2 earlier trials, 1 in Indonesia⁷ and another in south India,⁸ in which newborns dosed similarly also experienced reductions in mortality during the first months of life. On the other hand, these encouraging findings have not been duplicated in Africa. Perinatal VAS in a recent trial in Guinea-Bissau failed to achieve a benefit in mortality.⁹ An earlier randomized trial in Zimbabwe also showed no survival advantage for infants given VAS at birth.¹⁰

Various factors are involved that could confound the evidence supporting the benefit of VAS in neonates. For example, there is controversy about the effects that VAS has on the response to vaccines when given concomitantly with immunizations between the ages of 1 and 5 months. Insufficient data are currently available to demonstrate the effects of single-dose neonatal VAS on response to subsequent vaccinations.¹¹ The prevalence of vitamin A deficiency in mothers and their infants varies greatly between regions, which has yielded differential responses to neonatal supplementation programs. A high prevalence of diseases including HIV, measles, and

other infections may affect the efficacy of neonatal programs. Humphrey et al reported that VAS given at birth to either mothers or their HIV-exposed infants in Zimbabwe who were polymerase chain reaction–negative at 6 weeks doubled mortality by 2 years of age.¹² However, there was a mortality benefit in infants given VAS at birth who were HIV-positive at baseline. Genetic factors that may affect response to VAS have been identified in both human¹³ and animal¹⁴ studies. Benn et al¹⁵ have pointed out that boys seem to derive greater benefit from VAS than girls. This may be a result of greater vitamin A deficiency in boys, gender-related immunologic differences, or differential response to vaccines administered with VAS. It is worth noting that in the study by Klemm et al an opposing trend was reported: female infants seemed to benefit more than the male infants.¹

Klemm et al conducted a large, rigorous study in Bangladesh under difficult conditions. Given the lack of efficacy of VAS seen in infants aged 1 to 5 months in earlier studies, it seems that giving the supplementation close to birth is important. An open question is whether the level of public health coordination required to provide VAS within the first 24 hours of life is achievable in resource-poor countries, where most births occur at home. An encouraging sign in Bangladesh is the fact that since the 1980s the rate of BCG vaccination, targeted to be given at birth, rose from 2% to 95% by 2000,¹⁶ which raises hope that VAS and BCG vaccine could be administered concurrently after delivery.

In the Klemm et al study, an interesting observation was that there was a positive benefit even for those infants who were dosed nearly 1 month after birth. This finding needs to be understood in light of earlier studies that showed no benefit in VAS administered with vaccinations from the ages of 1 to 5 months. The effects of single versus multiple dosing, immunization interactions, susceptibility to infection, and age-dependent response to VAS all could be involved in explaining these different results. Although there was a benefit seen in infants who were receiving VAS even if their mothers

Abbreviation: VAS, vitamin A supplementation

Opinions expressed in these commentaries are those of the author and not necessarily those of the American Academy of Pediatrics or its Committees.

www.pediatrics.org/cgi/doi/10.1542/peds.2008-0455

doi:10.1542/peds.2008-0455

Accepted for publication Feb 19, 2008

Address correspondence to Steven A. Abrams, MD, Baylor College of Medicine, 1100 Bates St, Houston TX 77030. E-mail: sabrams@bcm.edu
 PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275). Copyright © 2008 by the American Academy of Pediatrics

were enrolled in the placebo arm of a concomitant weekly, low-dose VAS trial during pregnancy, it would be instructive to know if additional maternal factors, such as the effects of peripartum maternal retinol, affected infant survival.

It is clear that we have much to understand about neonatal VAS. In southern Asia, which has a relatively low overall rate of HIV infection compared with sub-Saharan Africa, there is a mortality benefit. It seems appropriate to move ahead with implementation of neonatal VAS as a public health policy in regions where it has been shown to have a survival benefit. However, implementation of programs should not proceed in any specific geographic area until controlled trials have shown a benefit in that region. Research that elucidates the underlying mechanisms of VAS and individual factors that may modify its benefit, such as gender, interaction with immunizations, coinfections, or exposure to diseases such as HIV and genetic and maternal factors, needs to continue. The goal should be to understand the effects and mechanisms of VAS adequately so that infants can be chosen for or excluded from supplementation on the basis of a defined set of criteria.

Neonatal VAS seems to be an intervention that holds promise in accomplishing a step-wise reduction in infant mortality in some resource-poor regions of the world, where successes have been difficult to achieve. However, we agree with others that caution and additional trials are needed before global use of such a program is undertaken.¹⁷

REFERENCES

1. Klemm RDW, Labrique AB, Christian P, et al. Newborn vitamin A supplementation reduced infant mortality in rural Bangladesh. *Pediatrics*. 2008;122(1). Available at: www.pediatrics.org/cgi/content/full/122/1/e242
2. Sommer A, Tarwotjo I, Djunaedi E, et al. Impact of vitamin A supplementation on childhood mortality: a randomised controlled community trial. *Lancet*. 1986;1(8491):1169–1173
3. Beaton GH, Martorell R, Aronson KJ, et al. *Effectiveness of Vitamin A Supplementation in the Control of Young Child Morbidity and Mortality in Developing Countries*. Geneva, Switzerland: World Health Organization; 1993. Standing Committee on Nutrition of the United Nations Administrative Committee (ACC/SCN) State of the Art Series Nutrition Policy discussion paper 13
4. Ching P, Birmingham M, Goodman T, Sutter R, Loevinsohn B. Childhood mortality impact and costs of integrating vitamin A supplementation into immunization campaigns. *Am J Pub Health*. 2000;90(10):1526–1529
5. Darlow BA, Graham PJ. Vitamin A supplementation to prevent mortality and short and long-term morbidity in very low birth-weight infants. *Cochrane Database Syst Rev*. 2007;(4):CD000501
6. West KP Jr, Katz J, Shrestha SR, et al. Mortality of infants <6 mo of age supplemented with vitamin A: a randomized, double-masked trial in Nepal. *Am J Clin Nutr*. 1995;62(1):143–148
7. Humphrey JH, Agoestina T, Wu L, et al. Impact of neonatal vitamin A supplementation on infant morbidity and mortality. *J Pediatr*. 1996;128(4):489–496
8. Rahmathullah L, Tielsch JM, Thulasiraj RD, et al. Impact of supplementing newborn infants with vitamin A on early infant mortality: community based randomised trial in southern India. *BMJ*. 2003;327(7409):254–257
9. Fisker AB, Lisse IM, Aaby P, et al. Effect of vitamin A supplementation with BCG vaccine at birth on vitamin A status at 6 wk and 4 mo of age. *Am J Clin Nutr*. 2007;86(4):1032–1039
10. Malaba LC, Iliff PJ, Nathoo KJ, et al. Effect of postpartum maternal or neonatal vitamin A supplementation on infant mortality among infants born to HIV-negative mothers in Zimbabwe. *Am J Clin Nutr*. 2005;81(2):454–460
11. Stephenson CB. Commentary: a hypothesis concerning vitamin A supplementation, vaccines, and childhood mortality. *Int J Epidemiol*. 2003;32(5):828–829
12. Humphrey JH, Iliff PJ, Marinda ET, et al. Effects of a single large dose of vitamin A, given during the postpartum period to HIV-positive women and their infants, on child HIV infection, HIV-free survival, and mortality. *J Infect Dis*. 2006;193(6):860–871
13. Kuhn L, Coutsooudis A, Trabatonni D, et al. Synergy between mannose-binding lectin gene polymorphisms and supplementation with vitamin A influences susceptibility to HIV infection in infants born to HIV-positive mothers. *Am J Clin Nutr*. 2006;84(3):610–615
14. Bhat PV. Retinal dehydrogenase gene expression in stomach and small intestine of rats during postnatal development and in vitamin A deficiency. *FEBS Lett*. 1998;426(2):260–262
15. Benn CS, Fisker AB, Diness BR, Aaby P. Neonatal vitamin A supplementation: sex-differential effects on mortality? *J Infect Dis*. 2006;194(5):719
16. Steele F, Diamond I, Amin S. Immunization uptake in rural Bangladesh: a multilevel analysis. *J R Stat Soc Ser A Stat Soc*. 1996;159(2):289–299
17. Fawzi WW. The benefits and concerns related to vitamin A supplementation. *J Infect Dis*. 2006;193(6):756–759

Postnatal Vitamin A Supplementation in Developing Countries: An Intervention Whose Time Has Come?

Steven A. Abrams and David C. Hilmers

Pediatrics 2008;122;180-181

DOI: 10.1542/peds.2008-0455

Updated Information & Services	including high-resolution figures, can be found at: http://www.pediatrics.org/cgi/content/full/122/1/180
References	This article cites 15 articles, 7 of which you can access for free at: http://www.pediatrics.org/cgi/content/full/122/1/180#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Nutrition & Metabolism http://www.pediatrics.org/cgi/collection/nutrition_and_metabolism
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.pediatrics.org/misc/Permissions.shtml
Reprints	Information about ordering reprints can be found online: http://www.pediatrics.org/misc/reprints.shtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

